

Drug Combo Improves Quality of Life for People With Liver Cancer

Tecentriq plus Avastin improved both overall survival and patient-reported outcomes.

January 26, 2020 By [Liz Highleyman](#)

A combination of the immune checkpoint inhibitor Tecentriq (atezolizumab) and the targeted therapy Avastin (bevacizumab) improved quality of life and delayed worsening of symptoms in people with advanced liver cancer, according to a report presented Friday at the American Society of Clinical Oncology (ASCO) Gastrointestinal Cancers Symposium in San Francisco.

The patient experience with cancer treatment was a key theme of this year's meeting, as the field of oncology increasingly recognizes the importance of not only extending life, but also ensuring that its quality is maintained.

"Patient-reported outcomes, or PROs, are now recognized as important endpoints of cancer clinical trials that provide important insights regarding the impact of treatment on patients' quality of life," ASCO chief medical officer and executive vice president Richard Schilsky, MD, said in a [press release](#). "PROs inform us about the tolerability of new therapies, which is just as important as efficacy in gauging their utility and acceptance by patients."

Over years or decades, chronic hepatitis B or C, heavy alcohol use, fatty liver disease and other causes can lead to the development of liver cirrhosis and hepatocellular carcinoma (HCC), the most common type of liver cancer. HCC is often detected late and is difficult to treat, making it one of the leading causes of cancer death.

Peter Galle, MD, PhD, of the University Medical Center in Mainz, Germany presented patient-reported outcomes from the Phase III IMbrave150 trial, which compared Tecentriq plus Avastin versus standard treatment using the targeted therapy Nexavar (sorafenib).

Tecentriq is a PD-L1 checkpoint inhibitor that restores T-cell activity against cancer. Avastin is a monoclonal antibody that blocks VEGF, a protein that promotes the proliferation of blood vessels that supply tumors and plays a role in immune suppression.

"Because it reflects both the effects of disease and the side effects of treatment, sustained or improved quality of life is particularly important for patients," Galle said. "Patients with liver cancer are typically more fragile and frail than others. Toxicity of the treatments can be much

more serious for these patients, and their quality of life can decline quite quickly.”

IMbrave included 501 people with inoperable locally advanced or metastatic liver cancer who had not received prior systemic treatment. They were randomly assigned to take Tecentriq plus Avastin, both administered by IV infusion every three weeks, or twice-daily oral Nexavar until they experienced disease progression or unacceptable side effects.

Primary results from the study were presented at the European Society for Medical Oncology Asia Congress in November, showing that Tecentriq plus Avastin reduced the risk of disease progression or death by 41% (median 6.8 versus 4.3 months). Overall survival was 13.2 months in the Nexavar group but was not reached in the combination group because most patients were still alive. Overall response rates, meaning complete or partial tumor shrinkage, were 27% versus 12%, respectively.

Looking at side effects, people taking Tecentriq plus Avastin were more likely to experience fever, abnormal liver and kidney biomarkers and infusion reactions, while those taking Nexavar had more diarrhea and hand-foot syndrome (palmar-plantar erythrodysesthesia), with redness, swelling and pain on the palms of the hands and soles of the feet.

At the recent conference, Galle presented details about patient experiences during the trial. Participants were asked to complete questionnaires every three weeks during treatment and every three months after discontinuation for a year. The surveys asked about quality of life, physical functioning and role functioning, or how well they were able to perform their usual social and occupational activities. Clinically meaningful deterioration was defined as a decrease of 10 or more points from baseline on a 100-point scale. More than 90% of people in both treatment groups completed the questionnaires through most of the treatment period.

Tecentriq plus Avastin recipients reported about a 30% decline in their quality of life, compared with about a 40% decline among Nexavar recipients. What’s more, the time to deterioration was substantially longer for those taking the combination (11.2 versus 3.6 months, respectively).

Physical functioning showed a similar pattern. Tecentriq plus Avastin recipients reported about a 23% decline versus about 37% among Nexavar recipients. Here too, the time to deterioration was longer in the former group (13.1 versus 4.9 months, respectively). The time to a decline in role functioning was also longer in the former group (9.1 versus 3.6 months, respectively).

Finally, Tecentriq plus Avastin recipients reported a longer period before they experienced worsening of symptoms including loss of appetite, diarrhea, fatigue and pain.

“High-quality patient-reported outcome data demonstrate clinical meaningful benefits in key aspects of the patient experience (quality of life, functioning, key symptoms) with atezolizumab plus bevacizumab versus sorafenib,” the researchers concluded.

Editor’s note: On January 27, [Genentech announced](#) that it has submitted a supplemental biologics

license application to the Food and Drug Administration for the approval of Tecentriq plus Avastin for people with unresectable HCC.

[Click here](#) to read the study abstract.

[Click here](#) to learn more about liver cancer.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.cancerhealth.com/article/drug-combo-improves-quality-life-people-liver-cancer>