

Many U.S. Families of Children With Leukemia May Experience Food Insecurity

What's more, receiving SNAP benefits was not enough to address food insecurity for many families of children with leukemia, a study found.

October 20, 2022 By American Association for Cancer Research

Nearly one in four families of pediatric patients with acute lymphoblastic leukemia (ALL) enrolled in a clinical trial experienced food insecurity, and almost half of the families eligible for the Supplemental Nutrition Assistance Program (SNAP) did not receive benefits. In addition, receiving SNAP benefits was not sufficient to address food insecurity, according to results presented at the [15th AACR \(American Association for Cancer Research\) Conference on the Science of Cancer Health Disparities in Racial/Ethnic Minorities and the Medically Underserved](#), held September 16 to 19.

“Food insecurity is connected to worse health outcomes in general pediatrics,” said Rahela Aziz-Bose, MD, a pediatric oncology fellow at Boston Children’s Hospital and Dana-Farber Cancer Institute and lead author of the study. “It is incredibly important to understand whether patients who are being treated for cancer are experiencing food insecurity, because it is a risk factor on which we can intervene, and we can potentially improve outcomes.”

Receipt of SNAP benefits has been shown to reduce food insecurity in general pediatrics, representing a resource to mitigate adverse social determinants of health for children. Aziz-Bose and colleagues analyzed the prevalence of low income and food insecurity among a group of patients with ALL, the most common childhood cancer, enrolled in the Dana-Farber Cancer Institute (DFCI)/ALL Consortium [clinical trial](#) at six U.S. cancer centers from 2017 to 2022. An ancillary study within this pediatric oncology trial prospectively collected parent-reported information on income, SNAP receipt, and food insecurity.

“We looked at how well these six cancer centers are doing in connecting these families to nutrition supports that already exist, like SNAP,” said Aziz-Bose. “In addition, we wanted to understand whether receiving SNAP helps relieve food insecurity for these families.”

The researchers utilized parent-reported income at diagnosis and six months into therapy as a proxy to identify families that were SNAP-eligible. According to federal guidelines, households with

income lower than 130 percent of the Federal Poverty Level are eligible for SNAP. The authors applied statistical methods to study the association between SNAP eligibility, SNAP receipt, and parent-reported food insecurity.

The study revealed that 21 percent of 262 families evaluable at diagnosis and 25 percent of 223 families evaluable six months into therapy reported food insecurity. “The most important finding was that poverty and food insecurity are very common in this group of patients,” said Aziz-Bose.

At diagnosis, 20 percent of the families were SNAP-eligible. Of these, 60 percent reported food insecurity but only 53 percent were receiving SNAP benefits. A similar scenario was observed six months into therapy, when 28 percent of the families were SNAP-eligible, of whom 58 percent reported food insecurity and 58 percent were receiving SNAP.

“Nearly half of the eligible families were not receiving SNAP benefits, even though their children were being treated at highly resourced academic cancer centers,” said Aziz-Bose.

Among the 33 families that were SNAP-eligible at both time points, the fraction receiving SNAP benefits increased from 52 to 70 percent.

“It is promising that we could help some of the families who were eligible at diagnosis to enroll and receive benefits,” said Aziz-Bose. “However, there were families who became newly eligible between the two time points and were not receiving benefits at six months, highlighting that poverty is a dynamic state—families can move from one category to another as they progress through treatment, especially as the effects of out-of-pocket payments, transportation to visits, and lost income from a parent/caregiver having to give up a job all add up.”

Importantly, the study also showed that receiving SNAP benefits was not enough to address food insecurity, as more than half of the families receiving benefits were still experiencing food insecurity both at diagnosis and at six months into therapy (61 percent and 56 percent, respectively.)

“We need to universally screen for financial hardship at multiple points throughout pediatric cancer treatment,” added Aziz-Bose. “Connecting eligible families to available benefits is an important first step, but our findings suggest that it might not be enough. We need to address food insecurity through multiple different avenues. Our study also suggests that because of the financial stress associated with cancer treatment, the SNAP benefit amount needs to be higher for these families.”

According to the authors, the main limitation of this study is that SNAP eligibility could only be assessed through a proxy measure (self-reported income) and, since this was a retrospective analysis, it could not be confirmed. In addition, a more detailed analysis is needed to determine the cause/effect relationship between food insecurity and receiving SNAP benefits.

This study was funded by the National Institutes of Health. Aziz-Bose reports no conflicts of interest.

This [news release](#) was published by the American Association for Cancer Research (AARC) on September 16, 2022.

© 2026 Smart + Strong All Rights Reserved.

<http://beta.docker.cancerhealth.com/article/many-us-families-children-leukemia-may-experience-food-insecurity>